### Upside down and backwards

Kyle Gannon

BICMR, Peking University kgannon@bicmr.pku.edu.cn

 $\Delta_{24}$  Workshop at Inner Mongolia University September 6, 2025

Joint work with Tomasz Rzepecki

A large part of the model theory research involving topological dynamics focuses on the space of global types which are finitely satisfiable in a fixed small model.

- A large part of the model theory research involving topological dynamics focuses on the space of global types which are finitely satisfiable in a fixed small model.
- This is a seemingly appropriate setting because this space of types, equipped with the Newelski product, is canonically isomorphic to the Ellis semigoup of a particular group action.

- A large part of the model theory research involving topological dynamics focuses on the space of global types which are finitely satisfiable in a fixed small model.
- This is a seemingly appropriate setting because this space of types, equipped with the Newelski product, is canonically isomorphic to the Ellis semigoup of a particular group action.
- However, the Newelski product naturally extends to the space of global types which are invariant over a fixed small model. Similarly, this space of types with the extended product forms a left-continuous compact Hausdorff semigroup and is thus also susceptible to *Ellis theory analysis*.

- A large part of the model theory research involving topological dynamics focuses on the space of global types which are finitely satisfiable in a fixed small model.
- This is a seemingly appropriate setting because this space of types, equipped with the Newelski product, is canonically isomorphic to the Ellis semigoup of a particular group action.
- However, the Newelski product naturally extends to the space of global types which are invariant over a fixed small model. Similarly, this space of types with the extended product forms a left-continuous compact Hausdorff semigroup and is thus also susceptible to *Ellis theory analysis*.
- We take this extended product seriously and study the minimal ideals and Ellis subgroups of the semigroup of invariant types as well as their connections to the semigroup of finitely satisfiable types.

- A large part of the model theory research involving topological dynamics focuses on the space of global types which are finitely satisfiable in a fixed small model.
- This is a seemingly appropriate setting because this space of types, equipped with the Newelski product, is canonically isomorphic to the Ellis semigoup of a particular group action.
- However, the Newelski product naturally extends to the space of global types which are invariant over a fixed small model. Similarly, this space of types with the extended product forms a left-continuous compact Hausdorff semigroup and is thus also susceptible to Ellis theory analysis.
- We take this extended product seriously and study the minimal ideals and Ellis subgroups of the semigroup of invariant types as well as their connections to the semigroup of finitely satisfiable types.
- **5** We primarily focus on definably amenable NIP groups.

### Outline

Our talk is outlined as follows:

- Semigroup theory
- Model theoretic dynamics
- New results

Semigroups arise naturally in the study of dynamical.

Semigroups arise naturally in the study of dynamical. We recall some terminology from the theory of semigroups.

Semigroups arise naturally in the study of dynamical. We recall some terminology from the theory of semigroups.

Suppose that Y is a compact Hausdorff spaces and (Y, \*) is a semigroup.

Semigroups arise naturally in the study of dynamical. We recall some terminology from the theory of semigroups.

Suppose that Y is a compact Hausdorff spaces and (Y, \*) is a semigroup.

**3** A subset  $I \subseteq Y$  is called a **left ideal** of Y if I is closed and  $Y * I \subseteq I$ . We say that I is a **minimal left ideal** if I is a left ideal and does not properly contain any other left ideal.

Semigroups arise naturally in the study of dynamical. We recall some terminology from the theory of semigroups.

Suppose that Y is a compact Hausdorff spaces and (Y, \*) is a semigroup.

- A subset I ⊆ Y is called a left ideal of Y if I is closed and Y \* I ⊆ I. We say that I is a minimal left ideal if I is a left ideal and does not properly contain any other left ideal.
- ② We say that (Y, \*) is **left-continuous** if for every  $p \in Y$  the map  $-*p: Y \to Y$  is continuous.

Semigroups arise naturally in the study of dynamical. We recall some terminology from the theory of semigroups.

Suppose that Y is a compact Hausdorff spaces and (Y, \*) is a semigroup.

- A subset I ⊆ Y is called a left ideal of Y if I is closed and Y \* I ⊆ I. We say that I is a minimal left ideal if I is a left ideal and does not properly contain any other left ideal.
- ② We say that (Y, \*) is **left-continuous** if for every  $p \in Y$  the map  $-*p: Y \to Y$  is continuous.
- **3** Let  $u \in Y$ . We say that u is **idempotent** if u \* u = u.

Semigroups arise naturally in the study of dynamical. We recall some terminology from the theory of semigroups.

Suppose that Y is a compact Hausdorff spaces and (Y, \*) is a semigroup.

- A subset I ⊆ Y is called a left ideal of Y if I is closed and Y \* I ⊆ I. We say that I is a minimal left ideal if I is a left ideal and does not properly contain any other left ideal.
- ② We say that (Y, \*) is **left-continuous** if for every  $p \in Y$  the map  $-*p: Y \to Y$  is continuous.
- **3** Let  $u \in Y$ . We say that u is **idempotent** if u \* u = u.

The minimal left ideals of left-continuous compact Hausdorff semigroups admit a strong decomposition theorem.

#### Theorem (Ellis, Nerurkar)

Suppose that (Y,\*) is a left-continuous compact Hausdorff semigroup. Then (Y,\*) admits a minimal left ideal.

#### Theorem (Ellis, Nerurkar)

### Theorem (Ellis, Nerurkar)

Suppose that (Y,\*) is a left-continuous compact Hausdorff semigroup. Then (Y,\*) admits a minimal left ideal. Let I be a minimal left ideal of Y.

**1** There exists some  $u \in I$  such that u is idempotent, i.e. u \* u = u.

### Theorem (Ellis, Nerurkar)

- There exists some  $u \in I$  such that u is idempotent, i.e. u \* u = u.
- ② For each idempotent  $u \in I$ , u \* I is a group.


### Theorem (Ellis, Nerurkar)

- There exists some  $u \in I$  such that u is idempotent, i.e. u \* u = u.
- ② For each idempotent  $u \in I$ , u \* I is a group.
- **③** For any minimal left ideal J and idempotent  $v \in J$  we have that v \* J is isomorphic to u \* I (as an abstract group). We call v \* J an **Ellis** subgroup of Y.

#### Theorem (Ellis, Nerurkar)

- **1** There exists some  $u \in I$  such that u is idempotent, i.e. u \* u = u.
- ② For each idempotent  $u \in I$ , u \* I is a group.
- **③** For any minimal left ideal J and idempotent  $v \in J$  we have that v \* J is isomorphic to u \* I (as an abstract group). We call v \* J an **Ellis** subgroup of Y.
- I is the disjoint union of its Ellis subgroups. More explicitly, if id(I) is the collection of idempotents in I, then

$$I = \bigsqcup_{u \in \mathrm{id}(I)} u * I.$$







### Take-away

Take away: Given a left continuous compact Hausdorff semigroup, one can associate a canonical group object, i.e., the isomorphism type of the Ellis subgroups.

Question: Where do these kinds of semigroups arise in model theory?

Question: Where do these kinds of semigroups arise in model theory? Over certain types spaces with the Newelski product.

Question: Where do these kinds of semigroups arise in model theory? Over certain types spaces with the Newelski product.

Let G be a group, so a structure in a language  $\mathcal{L} := \{\cdot, 1, ^{-1}, ...\}$  which models the axioms of a group. Let  $\mathcal{G}$  be a monster model so that  $G \prec \mathcal{G}$ .

Question: Where do these kinds of semigroups arise in model theory? Over certain types spaces with the Newelski product.

Let G be a group, so a structure in a language  $\mathcal{L}:=\{\cdot,1,^{-1},...\}$  which models the axioms of a group. Let  $\mathcal{G}$  be a monster model so that  $G\prec\mathcal{G}$ .

**3** A type  $p \in S_x(\mathcal{G})$  is said to be invariant over G if whenever we have  $\varphi(x,b) \in p$  and  $a \equiv_G b$ , then  $\varphi(x,a) \in p$ .

Question: Where do these kinds of semigroups arise in model theory? Over certain types spaces with the Newelski product.

Let G be a group, so a structure in a language  $\mathcal{L} := \{\cdot, 1, ^{-1}, ...\}$  which models the axioms of a group. Let  $\mathcal{G}$  be a monster model so that  $G \prec \mathcal{G}$ .

- **1** A type  $p \in S_x(\mathcal{G})$  is said to be invariant over G if whenever we have  $\varphi(x,b) \in p$  and  $a \equiv_G b$ , then  $\varphi(x,a) \in p$ .
- **3** We let  $S_x^{\text{inv}}(\mathcal{G}, G)$  denote the space of global types which are invariant over G.

Question: Where do these kinds of semigroups arise in model theory? Over certain types spaces with the Newelski product.

Let G be a group, so a structure in a language  $\mathcal{L} := \{\cdot, 1, ^{-1}, ...\}$  which models the axioms of a group. Let  $\mathcal{G}$  be a monster model so that  $G \prec \mathcal{G}$ .

- **1** A type  $p \in S_x(\mathcal{G})$  is said to be invariant over G if whenever we have  $\varphi(x,b) \in p$  and  $a \equiv_G b$ , then  $\varphi(x,a) \in p$ .
- **3** We let  $S_x^{\text{inv}}(\mathcal{G}, G)$  denote the space of global types which are invariant over G.
- **3** A type  $p \in S_x(\mathcal{G})$  is said to be finitely satisfiable in G if whenever we have  $\varphi(x,b) \in p$ , there exists some  $c \in G$  such that  $\mathcal{G} \models \varphi(c,b)$ .

Question: Where do these kinds of semigroups arise in model theory? Over certain types spaces with the Newelski product.

Let G be a group, so a structure in a language  $\mathcal{L} := \{\cdot, 1, ^{-1}, ...\}$  which models the axioms of a group. Let  $\mathcal{G}$  be a monster model so that  $G \prec \mathcal{G}$ .

- **1** A type  $p \in S_x(\mathcal{G})$  is said to be invariant over G if whenever we have  $\varphi(x,b) \in p$  and  $a \equiv_G b$ , then  $\varphi(x,a) \in p$ .
- **3** We let  $S_x^{\text{inv}}(\mathcal{G}, G)$  denote the space of global types which are invariant over G.
- **3** A type  $p \in S_x(\mathcal{G})$  is said to be finitely satisfiable in G if whenever we have  $\varphi(x,b) \in p$ , there exists some  $c \in G$  such that  $\mathcal{G} \models \varphi(c,b)$ .
- We let  $S_x^{fs}(\mathcal{G}, \mathcal{G})$  denote the space of global types which are finitely satisfiable in G.

Exercise:  $S_x^{fs}(\mathcal{G}, G) \subseteq S_x^{inv}(\mathcal{G}, G)$ .

We now define the semigroup operation on the space of types:

We now define the semigroup operation on the space of types:

Suppose that  $p, q \in S_x^{\text{inv}}(\mathcal{G}, G)$ . Then we say that  $\varphi(x, c) \in p * q$  if and only if  $\mathcal{G} \models \varphi(a \cdot b, c)$  where  $b \models p|_{Gc}$ , and  $a \models p|_{Gcb}$ .

### Newelski Product

We now define the semigroup operation on the space of types:

Suppose that  $p, q \in S_x^{\text{inv}}(\mathcal{G}, G)$ . Then we say that  $\varphi(x, c) \in p * q$  if and only if  $\mathcal{G} \models \varphi(a \cdot b, c)$  where  $b \models p|_{Gc}$ , and  $a \models p|_{Gcb}$ .

Intuitively, realize q, realize p over the realization of q, then consider the type of the product.

#### **Folklore**

Both  $S_x^{\text{inv}}(\mathcal{G},G)$  and  $S_x^{\text{fs}}(\mathcal{G},G)$  with the Newelski product are left-continuous compact Hausdorff groups.

### Newelski Product

We now define the semigroup operation on the space of types:

Suppose that  $p, q \in S_x^{\text{inv}}(\mathcal{G}, G)$ . Then we say that  $\varphi(x, c) \in p * q$  if and only if  $\mathcal{G} \models \varphi(a \cdot b, c)$  where  $b \models p|_{Gc}$ , and  $a \models p|_{Gcb}$ .

Intuitively, realize q, realize p over the realization of q, then consider the type of the product.

#### **Folklore**

Both  $S_x^{\text{inv}}(\mathcal{G},G)$  and  $S_x^{\text{fs}}(\mathcal{G},G)$  with the Newelski product are left-continuous compact Hausdorff groups.

The space  $S_x^{fs}(\mathcal{G}, G)$  has been extensively studied. Why?

### Newelski Product

We now define the semigroup operation on the space of types:

Suppose that  $p,q \in S_x^{\text{inv}}(\mathcal{G},G)$ . Then we say that  $\varphi(x,c) \in p*q$  if and only if  $\mathcal{G} \models \varphi(a \cdot b,c)$  where  $b \models p|_{Gc}$ , and  $a \models p|_{Gcb}$ .

Intuitively, realize q, realize p over the realization of q, then consider the type of the product.

#### **Folklore**

Both  $S_x^{\text{inv}}(\mathcal{G}, G)$  and  $S_x^{\text{fs}}(\mathcal{G}, G)$  with the Newelski product are left-continuous compact Hausdorff groups.

The space  $S_x^{fs}(\mathcal{G}, \mathcal{G})$  has been extensively studied. Why?

- **1** Historical; Newelski originally considered  $S_x(G)$ , when G is stable; extended to all types definable; then  $S_{\text{ext}}(G)$ ...
- $\mathcal{S}_{x}^{fs}(\mathcal{G},G)$  is isomorphic to an object from classical topological dynamics. Namely, the Ellis semigroup of a particular group action.

Consider  $G = (\mathbb{Z}; +, 0, <)$ . Then  $S_x^{inv}(\mathcal{G}, G)$  looks like the following:

### Example: Integers

Consider  $G = (\mathbb{Z}; +, 0, <)$ . Then  $S_x^{\text{inv}}(\mathcal{G}, G)$  looks like the following:

$$\begin{cases}
S_{x}^{fs}(\xi,G) \\
\frac{1}{2} & \frac{1}{2}
\end{cases}$$

$$\begin{cases}
S_{x}^{fs}(\xi,G) \\
\vdots \\
P_{o} & \frac{1}{2}
\end{cases}$$

$$\begin{cases}
S_{x}^{fs}(\xi,G) \\
P$$

## Example: Integers cont.

$$\begin{cases}
\zeta_0 & t_0 \neq \frac{1}{\zeta_0} & \downarrow \zeta_0
\end{cases}$$

Recall we are interested in the ideal groups of  $S_x^{fs}(\mathcal{G},G)$  and  $S_x^{inv}(\mathcal{G},G)$ . The case of  $S_x^{fs}(\mathcal{G},G)$  has the following history.

Recall we are interested in the ideal groups of  $S_x^{fs}(\mathcal{G}, G)$  and  $S_x^{inv}(\mathcal{G}, G)$ . The case of  $S_x^{fs}(\mathcal{G}, G)$  has the following history.

• Shelah: If  $\mathcal{G}$  is NIP, then we can associated to  $\mathcal{G}$  a canonical (topological group), namely  $\mathcal{G}/\mathcal{G}^{00}$  [ $\mathcal{G}^{00}$  is the smallest type-definable subgroup of bounded index].

Recall we are interested in the ideal groups of  $S_x^{fs}(\mathcal{G}, G)$  and  $S_x^{inv}(\mathcal{G}, G)$ . The case of  $S_x^{fs}(\mathcal{G}, G)$  has the following history.

- **3** Shelah: If  $\mathcal{G}$  is NIP, then we can associated to  $\mathcal{G}$  a canonical (topological group), namely  $\mathcal{G}/\mathcal{G}^{00}$  [ $\mathcal{G}^{00}$  is the smallest type-definable subgroup of bounded index].
- ② Newelski Conjecture: If  $\mathcal G$  is NIP, then the Ellis subgroups of  $S^{\mathrm{fs}}_{\mathsf x}(\mathcal G,G)$  are isomorphic to  $\mathcal G/\mathcal G^{00}$ .

Recall we are interested in the ideal groups of  $S_x^{fs}(\mathcal{G}, \mathcal{G})$  and  $S_x^{inv}(\mathcal{G}, \mathcal{G})$ . The case of  $S_x^{fs}(\mathcal{G}, \mathcal{G})$  has the following history.

- **③** Shelah: If  $\mathcal{G}$  is NIP, then we can associated to  $\mathcal{G}$  a canonical (topological group), namely  $\mathcal{G}/\mathcal{G}^{00}$  [ $\mathcal{G}^{00}$  is the smallest type-definable subgroup of bounded index].
- ② Newelski Conjecture: If  $\mathcal{G}$  is NIP, then the Ellis subgroups of  $S_x^{\mathsf{fs}}(\mathcal{G}, \mathcal{G})$  are isomorphic to  $\mathcal{G}/\mathcal{G}^{00}$ .
- **③** Pillay-Penazi-Gismutulin:  $SL_2(\mathbb{R})$  is a counterexample.

Recall we are interested in the ideal groups of  $S_x^{fs}(\mathcal{G}, \mathcal{G})$  and  $S_x^{inv}(\mathcal{G}, \mathcal{G})$ . The case of  $S_x^{fs}(\mathcal{G}, \mathcal{G})$  has the following history.

- **3** Shelah: If  $\mathcal{G}$  is NIP, then we can associated to  $\mathcal{G}$  a canonical (topological group), namely  $\mathcal{G}/\mathcal{G}^{00}$  [ $\mathcal{G}^{00}$  is the smallest type-definable subgroup of bounded index].
- ② Newelski Conjecture: If  $\mathcal{G}$  is NIP, then the Ellis subgroups of  $S_x^{\mathsf{fs}}(\mathcal{G}, \mathcal{G})$  are isomorphic to  $\mathcal{G}/\mathcal{G}^{00}$ .
- **③** Pillay-Penazi-Gismutulin:  $SL_2(\mathbb{R})$  is a counterexample.
- $\ \ \,$  Chernikov-Simon: If  $\mathcal G$  is NIP + definably amenable, then Newelski's conjecture holds.

Recall we are interested in the ideal groups of  $S_x^{fs}(\mathcal{G}, G)$  and  $S_x^{inv}(\mathcal{G}, G)$ . The case of  $S_x^{fs}(\mathcal{G}, G)$  has the following history.

- **3** Shelah: If  $\mathcal{G}$  is NIP, then we can associated to  $\mathcal{G}$  a canonical (topological group), namely  $\mathcal{G}/\mathcal{G}^{00}$  [ $\mathcal{G}^{00}$  is the smallest type-definable subgroup of bounded index].
- ② Newelski Conjecture: If  $\mathcal G$  is NIP, then the Ellis subgroups of  $S^{\mathsf{fs}}_{\mathsf x}(\mathcal G, G)$  are isomorphic to  $\mathcal G/\mathcal G^{00}$ .
- **③** Pillay-Penazi-Gismutulin:  $SL_2(\mathbb{R})$  is a counterexample.
- $\hbox{ \small \textbf{0} Chernikov-Simon: If $\mathcal{G}$ is NIP $+$ definably amenable, then Newelski's conjecture holds.}$

More explicitly, the quotient map  $\pi: \mathcal{G} \to \mathcal{G}/\mathcal{G}^{00}$  extends to a map  $\hat{\pi}: \mathcal{S}_{\kappa}(\mathcal{G})$  and  $\pi|_{u*I}: u*I \to \mathcal{G}/\mathcal{G}^{00}$  is an algebraic isomorphism [where I is a minimal left ideal and u is an idempotent in I].

## Definably amenable groups

#### Fact

Suppose that T is NIP. Then the following are equivalent:

- G is definably amenable, i.e., G admits a left invariant measures on the collection of definable subsets.
- ②  $S_x^{\text{inv}}(\mathcal{G}, G)$  admits a left (right) strong f-generic, i.e., there exists some  $p \in S_x^{\text{inv}}(\mathcal{G}, G)$  such that every global left (right) translate of p is still an element of  $S_x^{\text{inv}}(\mathcal{G}, G)$ .

If G is NIP and definably amenable, we let  $\mathcal{F}_r$  be the collection of global right strong f-generics.

# Ideal groups in $S_x^{\text{inv}}(\mathcal{G},\overline{G})$

So what about that case of  $S_x^{inv}(\mathcal{G}, G)$ ?

### Theorem (G.-Rzepecki 2025+)

Suppose that G is NIP and definably amenable. Then

- $S_x^{\text{inv}}(\mathcal{G}, G)$  contains a unique minimal left ideal,  $\mathcal{F}_r$ . These are precisely the *right strong f-generics*.
- ② If  $u \in \mathcal{F}_r$  is an idempotent, then  $u * S_x^{\text{inv}}(\mathcal{G}, G)$  is an ideal group.
- **1** (HPP) The right stabilizer of any strong right f-generic is  $\mathcal{G}/\mathcal{G}^{00}$ .
- **③** For any idempotent  $u \in \mathcal{F}_r$ ,  $\pi|_{u*S_x^{\mathsf{inv}}(\mathcal{G},G)} : u*S_x^{\mathsf{inv}}(\mathcal{G},G) \to \mathcal{G}/\mathcal{G}^{00}$  is an algebraic isomorphism.

As consequence, we have that the ideal group of  $S_x^{\text{inv}}(\mathcal{G},G)$  and  $S_x^{\text{fs}}(\mathcal{G},G)$  are always isomorphic.

# Ideal groups in $S_x^{inv}(\mathcal{G}, \mathcal{G})$

So what about that case of  $S_x^{inv}(\mathcal{G}, \mathcal{G})$ ?

### Theorem (G.-Rzepecki 2025+)

Suppose that G is NIP and definably amenable. Then

- $S_x^{\text{inv}}(\mathcal{G}, G)$  contains a unique minimal left ideal,  $\mathcal{F}_r$ . These are precisely the *right strong f-generics*.
- ② If  $u \in \mathcal{F}_r$  is an idempotent, then  $u * S_x^{\text{inv}}(\mathcal{G}, G)$  is an ideal group.
- **1** (HPP) The right stabilizer of any strong right f-generic is  $\mathcal{G}/\mathcal{G}^{00}$ .
- **③** For any idempotent  $u \in \mathcal{F}_r$ ,  $\pi|_{u*S_x^{\mathsf{inv}}(\mathcal{G},G)} : u*S_x^{\mathsf{inv}}(\mathcal{G},G) \to \mathcal{G}/\mathcal{G}^{00}$  is an algebraic isomorphism.

As consequence, we have that the ideal group of  $S_x^{\rm inv}(\mathcal{G},G)$  and  $S_x^{\rm fs}(\mathcal{G},G)$  are always isomorphic. Furthermore, essentially the same map gives an isomorphism.

# Ideal groups in $S_x^{inv}(\mathcal{G}, \mathcal{G})$

So what about that case of  $S_x^{inv}(\mathcal{G}, \mathcal{G})$ ?

### Theorem (G.-Rzepecki 2025+)

Suppose that G is NIP and definably amenable. Then

- **3**  $S_x^{\text{inv}}(\mathcal{G}, \mathcal{G})$  contains a unique minimal left ideal,  $\mathcal{F}_r$ . These are precisely the *right strong f-generics*.
- ② If  $u \in \mathcal{F}_r$  is an idempotent, then  $u * S_x^{inv}(\mathcal{G}, G)$  is an ideal group.
- **1** (HPP) The right stabilizer of any strong right f-generic is  $\mathcal{G}/\mathcal{G}^{00}$ .
- For any idempotent  $u \in \mathcal{F}_r$ ,  $\pi|_{u*S_x^{\mathsf{inv}}(\mathcal{G},G)} : u*S_x^{\mathsf{inv}}(\mathcal{G},G) \to \mathcal{G}/\mathcal{G}^{00}$  is an algebraic isomorphism.

As consequence, we have that the ideal group of  $S^{\rm inv}_{x}(\mathcal{G},\mathcal{G})$  and  $S^{\rm fs}_{x}(\mathcal{G},\mathcal{G})$  are always isomorphic. Furthermore, essentially the same map gives an isomorphism.

Ok - So, is there some kind of natural isomorphism?

# Ideal groups in $S_x^{inv}(\mathcal{G}, \mathcal{G})$

So what about that case of  $S_x^{inv}(\mathcal{G}, G)$ ?

### Theorem (G.-Rzepecki 2025+)

Suppose that G is NIP and definably amenable. Then

- **3**  $S_x^{\text{inv}}(\mathcal{G}, \mathcal{G})$  contains a unique minimal left ideal,  $\mathcal{F}_r$ . These are precisely the *right strong f-generics*.
- ② If  $u \in \mathcal{F}_r$  is an idempotent, then  $u * S_x^{inv}(\mathcal{G}, G)$  is an ideal group.
- **1** (HPP) The right stabilizer of any strong right f-generic is  $\mathcal{G}/\mathcal{G}^{00}$ .
- For any idempotent  $u \in \mathcal{F}_r$ ,  $\pi|_{u*S_x^{\mathsf{inv}}(\mathcal{G},G)} : u*S_x^{\mathsf{inv}}(\mathcal{G},G) \to \mathcal{G}/\mathcal{G}^{00}$  is an algebraic isomorphism.

As consequence, we have that the ideal group of  $S^{\rm inv}_{x}(\mathcal{G},\mathcal{G})$  and  $S^{\rm fs}_{x}(\mathcal{G},\mathcal{G})$  are always isomorphic. Furthermore, essentially the same map gives an isomorphism.

Ok - So, is there some kind of natural isomorphism? (Reconsider  $\mathbb{Z}$ )

### Retraction

If the underlying theory is NIP, there exists a mysterious map

$$F: \mathcal{S}_{\scriptscriptstyle{X}}^{\mathsf{inv}}(\mathcal{G}, \mathcal{G}) 
ightarrow \mathcal{S}_{\scriptscriptstyle{X}}^{\mathsf{fs}}(\mathcal{G}, \mathcal{G})$$

whose definition is a little bit delicate.

### Fact (Simon)

The retraction map F from  $S_x^{\text{inv}}(\mathcal{G},G)$  to  $S_x^{\text{fs}}(\mathcal{G},G)$  has the following properties: Let  $p,q\in S_x^{\text{inv}}(\mathcal{G},G)$ , then

- F is continuous,
- **1** If p is finitely satisfiable in M, then F(p) = p,
- For any M-definable function f,  $f_*(F(p)) = F(f_*(p))$ .
- **1** If q is finitely satisfiable in M, then  $F(q_x \otimes p_y) = q_x \otimes F(p_y)$ .

Question: Does the retraction map induce an isomorphism between Ellis subgroups?

## It's complicated

Sometimes? It's a little complicated...

# Abelian groups

It is essentially true for abelian NIP groups.

### Abelian groups

It is essentially true for abelian NIP groups.

#### Lemma

Suppose that G is NIP and definably amenable. Let I be a minimal left ideal contained in  $F(\mathcal{F}_r)$ . Fix an idempotent  $u \in I$ . Consider  $t \in \mathcal{F}_r$  such that F(t) = u. Suppose that every coset of  $\mathcal{G}^{00}$  has a representative in G. Then the following are equivalent:

- **1** For every  $g \in G(M)$ ,  $u \cdot g \in u * I$ .
- ②  $F_M|_{t*S^{\text{inv}}_G(\mathcal{U},M)}: t*S^{\text{inv}}_G(\mathcal{U},M) \to u*I$  is an isomorphism of Ellis subgroups.

New results

## Abelian groups

It is essentially true for abelian NIP groups.

#### Lemma

Suppose that G is NIP and definably amenable. Let I be a minimal left ideal contained in  $F(\mathcal{F}_r)$ . Fix an idempotent  $u \in I$ . Consider  $t \in \mathcal{F}_r$  such that F(t) = u. Suppose that every coset of  $\mathcal{G}^{00}$  has a representative in G. Then the following are equivalent:

- For every  $g \in G(M)$ ,  $u \cdot g \in u * I$ .
- $② F_M|_{t*S^{\text{inv}}_G(\mathcal{U},M)}: t*S^{\text{inv}}_G(\mathcal{U},M) \to u*I \text{ is an isomorphism of Ellis subgroups}.$

#### Theorem

Suppose G is NIP, abelian, and G contains representatives for each coset of  $\mathcal{G}^{00}$ . Let I be a minimal left ideal of  $F(\mathcal{F}_r)$ . Fix an idempotent  $u \in I$ . Consider  $t \in \mathcal{F}_r$  such that  $F_M(t) = u$ . Then the map

 $F_M|_{t*S_G^{\text{inv}}(\mathcal{U},M)}: t*S_G^{\text{inv}}(\mathcal{U},M) \to u*I \text{ is an isomorphism of Ellis subgroups.}$ 

## Example


The retraction map is not an isomorphism of Ellis groups in  $\mathbb{R} \rtimes \{\pm 1\}$ .

### Example

The retraction map is not an isomorphism of Ellis groups in  $\mathbb{R} \rtimes \{\pm 1\}$ . Elements of  $\mathbb{R} \rtimes \{\pm 1\}$  are elements of the cartesian product  $\mathbb{R} \times \{\pm 1\}$  with the following group law:

$$(a,i)\cdot (b,j) = \begin{cases} (a+b,j) & \text{if } i=1, \\ (a-b,-j) & \text{if } i=-1. \end{cases}$$

## $\mathbb{R} times \{\pm 1\}$



\*  $H_1$ ,  $H_2$  are ideal Subgroups of  $S_x^{inv}(E,G)$ .  $K_1$ ,  $K_2$  are ideal Subgroups of  $S_x^{fs}(E,G)$ 

Consider the model theoretic inversion,  $^{-1}: S_x(\mathcal{G}) \to S_x(\mathcal{G})$  defined via  $p^{-1} = \operatorname{tp}(a^{-1}/\mathcal{G})$ .

Consider the model theoretic inversion,  $^{-1}:S_x(\mathcal{G})\to S_x(\mathcal{G})$  defined via  $p^{-1}=\operatorname{tp}(a^{-1}/\mathcal{G})$ .

We remark that the inversion map sends invariant types to invariant type, finitely satisfiable types to finitely satisfiable types.

Consider the model theoretic inversion,  $^{-1}: S_x(\mathcal{G}) \to S_x(\mathcal{G})$  defined via  $p^{-1} = \operatorname{tp}(a^{-1}/\mathcal{G})$ .

We remark that the inversion map sends invariant types to invariant type, finitely satisfiable types to finitely satisfiable types.

#### Definition

We say that a definably amenable G is dfg if there exists some  $p \in S_x^{\text{inv}}(\mathcal{G}, G)$  such that p is definable over G and every global translate of p is also definable over G.

Intuition: dfg groups are the opposite of compact.

Consider the model theoretic inversion,  $^{-1}: S_x(\mathcal{G}) \to S_x(\mathcal{G})$  defined via  $p^{-1} = \operatorname{tp}(a^{-1}/\mathcal{G})$ .

We remark that the inversion map sends invariant types to invariant type, finitely satisfiable types to finitely satisfiable types.

#### Definition

We say that a definably amenable G is dfg if there exists some  $p \in S_x^{\text{inv}}(\mathcal{G}, G)$  such that p is definable over G and every global translate of p is also definable over G.

Intuition: dfg groups are the opposite of compact.

### Theorem (G.-Rzepecki 2025+)

Suppose that T is NIP, G is dfg, and t is a right dfg type over M. Then  $F \circ^{-1}|_{t*S^{inv}_G(\mathcal{U},M)}$  is an anti-isomorphism from an invariant Ellis subgroup to a finitely satisfiable Ellis subgroup. Precomposing with group inversion give an honest-to-goodness isomrophism.

## Thank you

Thank you!